

# **Bleeding, Broken Bones** and Seizure

B. Bleeding Apply direct pressure

2. Call for emergency help.

C. Shock Keep the victim calm and

3. Do not try to push the broken

moist dressing to prevent drying

4. Do not try to straighten out a

fracture. Let a doctor or trained

6. Splint unstable fractures to

It is an alarming sight; a person

whose limbs jerk violently, whose

eves may roll upward, whose breath

may become heavy with dribbling or

even frothing at the mouth. Breathing

may stop in some seizures, or the

to force anything into the victim's

1. During the seizure:

victim may bite his or her tongue so

severely that it may bleed and cause

an airway obstruction. Do not attempt

mouth. You may injure the victim and

A. There is little you can do to stop

C. Let the seizure run its course. D. Help the victim to lie down and

keep from falling, to avoid injury.

G. Do not try to restrain a seizure

H. Move objects out of the way

which may injure the victim (i.e. desk, table, chair, etc.).

I. If an object endangers the victim and cannot be moved, put clothing or

soft material between the seizure victim and the object.

E. Loosen restrictive clothing.

prevent painful motion.

5. Do not permit the victim to walk

bone back into place if it is sticking out of the skin. Do apply a

over the site

warm.

out.

about.

Seizure

yourself.

the seizure.

B. Call for help.

F. Use no force.

victim.

person do that.

#### **Head Injuries**

If there is bleeding from an ear, it can mean that there is a skull fracture.

1. Special care must be taken when trying to stop any scalp bleeding when there is a suspected skull fracture. Bleeding from the scalp can be very heavy even when the injury is not too serious.

2. Don't press too hard. Be extremely careful when applying pressure over the wound so that bone chips from a possible fracture will not be pressed into the brain.

3. Always suspect a neck injury when there is a serious head injury. Immobilize the head and neck.

4. Call for emergency help. Let a professional medical person clean the wound and stitch it. if necessary.

5. Do not give alcohol, cigarettes or other drugs. They may mask important symptoms.

#### **Internal Bleeding**

Warning Signs: coughing or vomiting up blood or "coffee ground" material. Passing blood in urine or stool. Passing black tar-like bowel movements. All require immediate medical attention!

1. Have the victim lie flat and relax

2. Do not let the victim take any medication or fluids by mouth until seen by a doctor who permits it.

3. Obtain emergency medical help immediately.

#### **Broken Bones**

Broken bones usually do not kill. Do not move the victim, unless the victim is in immediate danger of further injury.

1. Check for:

A. Breathing Give Rescue Breathing if needed.

## 2. After the seizure:

A. Check to see if the victim is breathing...If he or she is not ... give Rescue Breathing at once. (See Rescue Breathing section.) B. Check to see if the victim is wearing a Medic Alert, or similar. bracelet, necklace, etc. It describes emergency medical requirements.



C. Check to see if the victim has any burns around the mouth. This would indicate poison.

3. The victim of a seizure or convulsion may be conscious, but confused and not talkative when the intense movement stops. Stav with the victim. Be certain that breathing continues. Then, when the victim seems able to move, get medical attention.



# Poisoning and Burns Clibna enoteS

## Poisonina

The home is loaded with poisons: Cosmetics, Detergents, Bleaches, Cleaning Solutions, Glue, Lye, Paint, Turpentine, Kerosene, Gasoline and other petroleum products. Alcohol. Aspirin and other medications, and on and on.

1. Small children are most often the victims of accidental poisoning. If a child has swallowed or is suspected to have swallowed any substance that might be poisonous, assume the worst-Take Action.

2. Call your Poison Control Center. If none is in your area, call your emergency medical rescue squad. Bring suspected item and container with you.

A. Do not give counteragents unless directed to by Poison Control Center or physician.

B. Do not follow directions for neutralizing poisons found on poison container

C. If victim is conscious, dilute poison by giving moderate amounts of water

3. What you can do if the victim is unconscious:

A. Make sure patient is breathing. If not, tilt head back and perform mouth to nose breathing. Do not give anything by mouth. Do not attempt to stimulate person. Call emergency rescue squad immediately.

side so that the person will not choke on what is brought up.



5. Be prepared. Determine and verify your Poison Control Center and Fire Department Rescue Squad numbers and keep them near your telephone.

Alcohol is as much a poison as stimulants, tranquilizers, narcotics, hallucinogens or inhalants. Don't take drunkenness lightly. Too much alcohol can kill.

and pulse. If breathing has stopped or is very weak give Rescue Breathing. Caution: Victims being revived of alcohol poisoning can be violent. Be careful! They can harm themselves and others.

3. While waiting for help:

C. Do not throw water on the victim's face.

D. Do not give liquor or a stimulant.



4. If the victim is vomiting:

A. Roll him or her over onto the left



Burns Flame Burns Cool with water to stop the burning process. Remove garments and jewelry and

### **Drug Overdose**

A drug overdose is a poisoning.

1. Call for emergency help at once.


2. Check the victim's breathing

A. Watch breathing. B. Cover the person with a blanket

for warmth.


Remember: alcohol in combination with certain other drugs can be deadly!

Chemical burns of the eye require



cover burn victim with clean sheets

Call for help immediately



immediate medical attention after flushing with water for 20 minutes.

Call for help immediately.



or towels

A45

444